

Integrative adaptive management to address interactions between biological invasions and protected area connectivity: a Canadian perspective

Stuart W. Livingstone (10°1), Josie Hughes (10°, Richard Pither (10°, and Marie-Josée Fortin (10°)

^aDepartment of Ecology and Evolutionary Biology, University of Toronto, ON, Canada; ^bDepartment of Physical and Environmental Sciences, University of Toronto-Scarborough, ON, Canada; ^cEnvironment and Climate Change Canada-Landscape Ecology Research Section, Ottawa, ON, Canada

Corresponding author: Stuart W. Livingstone (email: s.livingstone@utoronto.ca)

Abstract

Expanding and creating protected area networks has become a central pillar of global conservation planning. In the management and design of protected area networks, we must consider not only the positive aspects of landscape connectivity but also how that connectivity may facilitate the spread of invasive species, a challenge that has become known as the connectivity conundrum. Here, we review key considerations for landscape connectivity planning for protected area networks, focusing on interactions between network connectivity and the management of invasive species. We propose an integrative adaptive management framework for protected area network planning with five main elements, including monitoring, budgeting considerations, risk assessment, inter-organizational coordination, and local engagement. Protected area planners can address the dynamic aspects of the connectivity conundrum through collaborative and integrative adaptive management planning.

Key words: connectivity, conservation, protected areas, invasive species, adaptive management, terrestrial

Introduction

Spatial planning for terrestrial protected areas (hereafter PAs) has been of central importance to the discipline of conservation biology for decades (Noss and Harris 1986) and is increasingly important for guiding conservation decisions at global, regional, and national scales. Protected area planners are faced with the challenge of mitigating human environmental impacts, which continue to cause a steady decline in biodiversity and ecological integrity in and around PAs across the planet (Beyer et al. 2019). Addressing these impacts includes planning for PA ecological connectivity, which is now seen as vital for addressing global environmental change (Liang et al. 2018; Stewart et al. 2019). Connectivity amongst PAs facilitates critical ecological processes such as dispersal, seasonal migrations, and species range shifts resulting from climate change, and in doing so, it can prevent deleterious effects, such as inbreeding and local extinctions, thereby helping to maintain ecosystem integrity (Saura et al. 2019). However, ecological connectivity can also have negative effects such as facilitating the spread of disturbances and invasive species. For example, well-connected PAs can act as corridors for the movement of exotic invasive species, such as the emerald ash borer (Agrilus planipennis Fairmaire) or the European fire ant (Myrmica rubra L.) (Resasco et al. 2014; Cuddington et al. 2018). Outbreaks of native species such as mountain pine beetles (Dendroctonous ponderosae Hopkins; Maguire et al.

2015) and spruce budworms (*Choristoneura occidentalis* Freeman; Drever et al. 2018) can also be facilitated by large contiguous areas of mature coniferous forest. Moreover, the connectivity of anthropogenic disturbances (e.g., road and trail networks, energy infrastructure corridors, and agricultural landscapes) can also facilitate the movement of non-desirable species within and between PAs (Schulze et al. 2018). This can alter predator-prey interactions and facilitate disturbancemediated species invasions into PAs (Vardarman et al. 2018).

The recently developed United Nations Kunming-Montreal Global Biodiversity Framework reaffirmed targets for having well-connected PAs and for mitigating the spread and impacts of invasive species (UN Environment Program 2022; Targets 2/3 and 6, respectively, of Goal A: 15th Conference of the Parties to the Convention on Biological Diversity). To achieve these targets, PA planners are developing proactive strategies to address the multifaceted challenges of global environmental change (Kullberg et al. 2019; Hilty et al. 2020), including land-use change, climate change (D'Aloia et al. 2019; Hilty et al. 2020), and biological invasions (Schulze et al. 2018; Hilty et al. 2020). However, in planning for PA networks and connectivity, integrative management approaches that account for both the benefits and risks of connectivity will be required. In Canada, the adoption of the Kunming-Montreal Global Biodiversity Framework will lead to the creation of a new suite of national goals and targets for conservation that will be formalized in 2024 (Environment and Climate Change Canada 2023). To achieve the new global and domestic targets, Canada will likely build upon existing initiatives for PA connectivity planning (e.g., the National Program for Ecological Corridors; Parks Canada Agency 2022) and for the mitigation of species invasions (Environment and Climate Change Canada 2016; Beazley et al. 2023). Whether for existing or new initiatives, managers will increasingly need to address these targets in tandem (Beger et al. 2022).

The threat of invasive species in protected areas

The spread of exotic invasive species represents a significant threat to the ecological integrity of PAs around the globe (Schulze et al. 2018). It is well known that invasive species can cause significant reductions in the diversity of native ecosystems as well as impairment of ecosystem functioning (Pyšek et al. 2012). In many cases, landscape alteration or degradation can create opportunities for non-native species to establish and potentially spread throughout a region (Hierro et al. 2006). Conversely, intact ecosystems are typically more resistant to invasion (Beaury et al. 2019), but the degree of resistance is also contingent on the particular characteristics or traits of potential invasive species (Martin and Marks 2006). Indeed, the susceptibility of an ecosystem or PA to biological invasion is not a static property but rather a dynamic and context-dependent spatio-temporal process (Clark and Johnston 2011), demanding an adaptive management response. In Canada, research has been conducted to model the spread of invasive species, and while some local-scale spatial models for invasive species spread have been explicitly focused on PAs (Sy et al. 2009), regional-scale modelling efforts tend not to make explicit linkages to PA management. Furthermore, while invasive species management considerations are embedded into a multitude of policies and strategic frameworks across Canada—including some that focus on PAs (Meloche and Murphy 2006)-integration is lacking across scales and sectors (Smith et al. 2014).

Biological invasions are widely modelled and managed through a stage-based approach (Fig. 1), with optimal management actions recommended for each stage (Richardson et al. 2000). For stage i (pre-introduction), managers evaluate the potential pathways for invasion and act to minimize the possibility of transport. For stage *ii* (introduction and establishment), early detection and eradication are prioritized, often involving spatial niche modelling for specific species, which can inform monitoring efforts for vulnerable areas (Václavík et al. 2010). In stage iii (spread), the ability to contain spread and/or hinder dispersal is assessed (Mortensen et al. 2009). In stage iv (dominance and/or naturalization), the ability to reduce impact and prevent spread to other regions is prioritized, but such efforts are often confounded by a lack of coordination across jurisdictions, land tenure heterogeneity, and/or a lack of management resources (Epanchin-Niell et al. 2010). Managers also contend with the potential for wellconnected desirable habitat to allow for the spread of "native invasions" or "overabundant species" (Environment and Climate Change Canada 2010; Wilkerson 2013), which do not necessarily fit the stage-based model of invasion (Nackley et

al. 2017). Where non-native invasions tend to act synergistically with other human disturbances, hyper-abundance of native species often arises due to human environmental influence (e.g., resource subsidies, Lamarre et al. 2017) or can take the form of outbreaks (e.g., chronic wasting disease, Nobert et al. 2016; spruce budworm, Senf et al. 2017). These varied ecological possibilities present a major challenge for PA managers and the broader field of PA connectivity planning.

Variation in the movement of desirable and undesirable species across natural and anthropogenic landscape elements leads to a "connectivity conundrum", requiring PA planners to consider the positive and negative consequences of connectivity (Simberloff and Cox 1987; Hilty et al. 2020; Beger et al. 2022; Silva et al. 2023) and trade-offs in spatial conservation planning. Here, we elaborate on the confounding factors surrounding PA network planning and propose a framework to support such planning that recognizes variation in ecological interactions and the multifaceted nature of landscape connectivity. We argue for the importance of integrative adaptive management practices to address this complexity, including monitoring, human resource considerations, risk assessment, inter-organizational coordination, and local engagement. In the coming years in Canada, protected and conserved area planners will be developing and implementing new landscape connectivity and PA network projects and will simultaneously need to grapple with the management of invasive species. Below, we provide a synthetic perspective on these subjects that we hope can inform PA connectivity planning in Canada.

Confounding factors for protected area network planning

Anthropogenic contexts for protected area connectivity

The manner in which different forms and scales of human land-use activity affect biodiversity is highly variable (Decker et al. 2017). For example, intensive agriculture and/or urbanization typically have a direct negative effect on biodiversity (Newbold et al. 2015). These landscape processes tend to also be associated with the increased prevalence of non-native and invasive species (Cadotte et al. 2017), which often spill over into PAs (Padmanaba et al. 2017). Given that biological invasions can be facilitated by direct human transport or often act synergistically with other human disturbances, PAs located in areas with high-to-moderate human population density face the highest probability of invasion (Chapman et al. 2020). However, the presence of non-native invasive species in remote PAs has also been increasingly observed (Sanderson et al. 2012). Protected areas located in regions with high-tomoderate human population density tend to house a greater number of threatened species and generally have a high degree of biodiversity compared to more remote PAs (Kraus and Hebb 2020). These PAs are also differentially subjected to external pressures such as sound and light from anthropogenic sources and chemical pollution runoff, as well as internal pressures such as the transportation and hospitality infrastructure created to support PA visitation (Jones et al. 2018). In



Fig. 1. Protected area invasion scenarios based on different forms of landscape connectivity and disturbance. The top panel depicts the variable nature of invasion risk associated with the positioning of an invasive species relative to a protected species. The middle panel depicts simplified connectivity-disturbance scenarios between two protected areas, noting also the potential invasion risk associated with each scenario. Scenario A illustrates a hypothetical low-connectivity/high-disturbance case where a disturbance-dependent invasive species is able to spread across a disturbed landscape (shown as hatched lines) and invade a protected area. Scenario B shows the case where an adjoining road between two protected areas creates a high degree of both anthropogenic connectivity and disturbance, facilitating the spread of invasive species via road transportation (e.g., stowaways or intentional transport) as well as through road edges (i.e., establishment and spread opportunities for invasive species). Scenario C depicts a low-connectivity/low-disturbance scenario where a disturbance-dependent invader is subject to resistance across the landscape through an intact ecosystem in the matrix, the protected area edge, and/or the protected area itself (i.e., "diversity-resistance"). Scenario D depicts a high-connectivity/low-disturbance scenario where a corridor of desirable habitat may also facilitate the spread of invaders that are not dependent on disturbance for establishment. The bottom-left panel depicts the stage-based conceptualization of the invasion process (i: pre-introduction, ii: establishment, iii: spread, iv: dominance) and the invasion curve. Invasive species can be present at different stages of the invasion process in the matrix, the area surrounding a protected area, or within the protected area itself. Each of these possibilities differentially informs focal management action. The bottom-right panel notes some of the primary factors considered in management prioritization and decision support modelling.



intensively farmed or urbanized landscapes, the availability of desirable habitat is extremely reduced (Wilson et al. 2016). Remnant habitat patches can act as stepping stones, both for species of conservation interest and/or invasive species, as they disperse or are transported between larger and more intact habitats or PAs (Saura et al. 2014). Conversely, fragmentation that impedes the movement of desired species can also reduce the spread of an invasive species that has specific habitat requirements (e.g., apple snail, *Pomacea canaliculata* Lamarck, in wetland habitats; Pierre et al. 2017).

Transportation corridors in and around PAs can have a dual effect of reducing habitat connectivity and impeding the movement of some organisms (Trombulak and Frissell 2000), while also facilitating the spread and establishment of non-native invasive species by creating a well-connected disturbed landscape that many invasive species are able to exploit (With 2002; Riitters et al. 2018). Indeed, studies have documented how the connectivity and disturbed conditions created by road networks can facilitate the spread of invasive species beyond the direct vehicular transport of nonnative species (e.g., Muthukrishnan et al. 2018). These types of invasions are a product of the interaction between anthropogenic disturbance and connectivity across the landscape, where both factors are explicitly and spatially linked (Fig. 1). Collision mortality risks increase with traffic density, but the impact of smaller rural roads used for forestry, mining, and other resource extraction activities can also be substantial. Invasive species can be transported by users of rural roads into otherwise undisturbed habitats (Trombulak and Frissell 2000), and disturbed road edges create opportunities for disturbance exploiting invasives to thrive and spread along roads and into intact areas (e.g., Japanese knotweed, Fallopia japonica, Dauer and Jongejans 2013; Phragmites australis (Cav.) Trin. Ex Steud., Sciance et al. 2016). Resource roads also alter behaviour, movement, and mortality risks for native species such as wolves, bears, and woodland caribou, which can, in turn, alter trophic dynamics and population viability (Mumma et al. 2018; Dickie et al. 2020; Proctor et al. 2020; Whittington et al. 2022). Thus, even sparsely populated rural areas can be surprisingly altered. Resource roads and other resource extraction activities are not consistently mapped (Poley et al. 2022), so global, national, and continental analyses do not fully capture the intensity and extent of these disturbances. Even so, in some areas, large tracts of intact land do remain to facilitate the movement and migration of wide-ranging species and connectivity among PAs (Belote et al. 2017; Hirsh-Pearson et al. 2022; Hughes et al. 2023; Pither et al. 2023).

Variation in edge effects across protected areas and corridor linkages

Edge effects refer to the environmental conditions that arise at the boundary of different ecosystems or land-use types (Didham 2010). Typically, edge effects are assessed through a conservation lens as products of anthropogenic landscape fragmentation and disturbance (Harper and Macdonald 2002). The relative significance of edge effects in PA planning varies in relation to the type of ecosystem within a PA, specific conservation objectives (e.g., planning for species at risk, connectivity planning), as well as the composition of the matrix (Fig. 1). For instance, a PA consisting of mostly forest habitat that is surrounded by farmland or an urbanized landscape (e.g., Rouge National Urban Park in Toronto, Canada) will be subject to edge effects at the margins of intact forest habitat (e.g., due to variation in micro-climate, light regime, seed dispersal, and colonization; Matlack 1993). Many invasive plant or arthropod species possess traits that allow them to take advantage of edge effects, readily dispersing along forest edges and establishing populations (Dillon et al. 2018). But many of these species that are able to colonize such disturbed habitats are limited in their ability to spread to intact habitats (Foxcroft et al. 2011). Yet, it can also be the case that an invasive species is able to establish itself in a disturbed edge environment and then spread to intact habitats, which is also a concern for the creation of new ecological corridors where edges and early successional habitats can allow for the establishment and spread of invasive species into PAs (Wilkerson 2013). This can occur due to an invader occupying an empty niche in the intact ecosystem and perhaps also possessing a fitness advantage that allows them to outcompete native species and exert impacts on the system (MacDougall et al. 2009). Vincetoxicum rossicum (Kleopow) Babar., an invasive plant species in Rouge National Urban Park, is an example of an invader that has been able to colonize disturbed edge habitat and spread to intact habitat in the PA and throughout the broader region (Sodhi et al. 2019). Similarly, P. australis has colonized disturbed edge habitats across Canada-adjacent to both natural and anthropogenic landscape features-and has also spread and become dominant in areas of significant conservation value (Jung et al. 2017).

For PA connectivity considerations, the potentially deleterious impacts of edge effects are of concern when planning for the creation and/or restoration of stepping-stone habitat patches or conservation corridors in fragmented landscapes. In these cases, managers are interested in whether disturbed corridors or patch edges can facilitate the spread of invasive species to valuable conservation land. A conservation corridor may consist of a wide stretch of relatively intact habitat that connects larger PAs, in which case, edge effects may be of minimal concern. Alternatively, a corridor could also consist of a newly restored habitat (i.e., early successional), a patchy and fragmented landscape with multiple land-use types and habitats, or a combination of these (Yu et al. 2012). In the case of a new conservation corridor where a large amount of land will undergo ecological restoration, monitoring can be prioritized during the early stages of restoration because a young system is often most susceptible to invasion (Fig. 2, #1; Yannelli et al. 2017). It is often the case that invasive species removal is the first step of restoration projects (Perry et al. 2017), where such efforts aim to optimize resistance to invasion (Funk et al. 2008).

Fig. 2. Building on existing best practices in invasive species and protected area management, we propose an integrative adaptive management framework for protected area network planning that addresses the interaction between these two conservation objectives. The center image depicts the spatial context for a hypothetical PA network, including the PAs (nodes), corridors (links), the matrix, and the potential for edge effects.



An integrative adaptive management framework addressing interactions between biological invasions and protected area connectivity

Protected areas around the globe are already dealing with biodiversity decline caused by anthropogenic disturbances and invasive species (Jones et al. 2018; Leberger et al. 2020). To address these dynamic challenges, individual PAs typically use adaptive management frameworks, and we argue that these approaches can be applied to the planning and management of PA connectivity initiatives in a manner that addresses the connectivity conundrum. We propose a framework with five main elements, ranging from assessing the current ecological condition to budgeting and developing partnerships.

Assessing ecological conditions

Canadian PAs are often governed by an adaptive management approach, where the status of the PA and management effectiveness are evaluated using ecological indicators and quantitative targets and thresholds (Wright et al. 2017). The key component of adaptive management for PAs is that ecosystem monitoring is carried out on a regular basis so that values for indicators can be assessed in relation to previous years and/or baseline conditions (Fig. 2, #1). With respect to the introduction and spread of invasive species in PAs and PA networks, ecosystem monitoring within an adaptive management program can allow for the identification and possible eradication of recently established invasive species and can also allow managers to evaluate the effectiveness of control methods (Rout et al. 2017). When resources permit, PA managers may apply an active adaptive management approach to gauge the relative effectiveness of multiple types of interventions for the control of abundant invasive species (e.g., physical removal, chemical control, and biological control) (Giljohann et al. 2011) or to examine the potential of different types of connectivity (e.g., stepping stones, intact corridors, and restored habitat corridor) to minimize the rate of spread across a PA connectivity project (Travers et al. 2021).

Monitoring is perhaps the most fundamental aspect of invasive species management. Goals include detecting newly established invaders, characterizing invader distribution and rate of spread, and assessing the effectiveness of different control measures (Foxcroft et al. 2017). However, a systematic monitoring approach for invasive species management does incur a significant financial cost to conservation managers, but the cost of managing well-established invasive species can be crippling (Moodley et al. 2022). Early detection of invasive species via ecological monitoring also maximizes the potential for eradication (Rejmánek and Pitcairn 2002).

With respect to monitoring the establishment and spread of invasive species and/or overabundant native species (Environment and Climate Change Canada 2010) between PAs or from unprotected land into PAs, different management considerations arise from conditions in the matrix between PAs, in the area surrounding a PA, or within the PA itself (Fig. 1). In planning for PA networks and/or connectivity between PAs, managers are faced with multiple possible scenarios and scales of invasion risk. For instance, PA managers may need to assess the risk associated with a specific invasive species that has yet to establish and develop a plan for proactive management actions (i.e., stages *i* and *ii*: pre-introduction and establishment; Fig. 1). It could also be the case that an invasive species is newly established and spreading in either a nearby PA or in the surrounding matrix. In this case, managers may need to work with regional stakeholders to coordinate monitoring and/or the application of control measures (i.e., stage iii: spread, Figs. 1 and 2, #5; e.g., Downey et al. 2010). For example, P. australis has dispersed across much of eastern North America, becoming common in both PAs and landscape corridors (natural and anthropogenic). Despite emergent bio-control options for P. australis management (Blossey et al. 2020) and the existence of inter-organizational and inter-national task forces (Great Lakes Phragmites Collaborative n.d.), it is most likely the case that P. australis will remain a significant challenge for ecosystem managers well into the future (Quirion et al. 2018). In this case, faced with a triage scenario, ecosystem and PA managers across the region are prioritizing the conservation of threatened species habitat (Markle et al. 2018).

Inter-organizational planning

The multi-scale and inter-jurisdictional nature of PA connectivity planning has spurred collaborative planning to identify opportunities to improve connectivity for conservation (Lemieux et al. 2021). Inter-organizational collaboration is also a common characteristic of invasive species management and risk assessment (Emilson and Stastny 2019), but the complexity of the issue often hinders effective intervention (Fantle-Lepczyk et al. 2022). Nevertheless, there are clear but unrealized synergies between the organizational structures involved in PA connectivity planning and invasive species management (Fig. 2, #2 and 5). Addressing the connectivity conundrum will require inter-organizational coordination that integrates connectivity for conservation, assessment of invasion risk, and public engagement (Bixler et al. 2016). For example, in the case of PA connectivity planning, local-scale land-use and management practices, both at the "node" scale (Häkkilä et al. 2018) and the "link" scale (corridors and matrix: Newmark et al. 2023), influence the quality and integrity of a larger PA network (Fig. 2). Such interactions can sometimes be complicated by the existence of multiple forms of land tenure and/or conflicting land-use practices across a proposed connectivity corridor or PA network (Mansourian et al. 2019; Hilty et al. 2020). These governance challenges that emerge in multi-actor and multi-scale interactions for PA planning are also pervasive in the practice of invasive species management (Estévez et al. 2015). However, the national-scale adoption and implementation of the Post-2020 Global Biodiversity Framework should promote data sharing and potentially spur the development of a national invasive species database (UN Environment Program, 2023; Fig. 2, #2 and 5), as has been proposed for the United States (Wallace et al. 2020). In Canada, organizations such as the Invasive Species Centre, the Canadian Council on Invasive Species, the National Indigenous Guardians Network, and a multitude of other regional partnerships are well positioned to establish working groups but are also faced with the hurdle of inadequate resource availability (Canadian Council on Invasive Species 2023). The integration of invasive species risk assessment with PA connectivity planning can provide a necessary proactive perspective to guide restoration efforts, land securement, and ecosystem monitoring.

Risk assessment and management prioritization

Effectively integrating invasive species risk assessment into PA connectivity planning initiatives will benefit from the consideration of predictive invasive species distribution models (SDMs). Species distribution models are widely used in conservation and ecology to predict biological responses to future environments (Lawler et al. 2011). Many SDM analyses are focused on species of conservation concern (Austin 2007), but there is an increasing interest in potential future distributions of invasive species (Srivastava 2019). For connectivity planning, SDMs can be developed to predict the risk of invasion into and across potential connectivity corridors by integrating a suite of ecological parameters (Stewart-Koster et al. 2015; Urziceanu et al. 2022). Species distribution models can also be synthesized with risk assessment frameworks to guide proactive monitoring efforts and management intervention strategies (Booy et al. 2017; Srivastava 2019). Some management plans account for the invasion stage, the spatial distribution of an invader, and dispersal predictions that may improve management effectiveness (Fournier and Turgeon 2017), sometimes emphasizing the probability of eradication to prioritize the timing of intervention efforts (Booy et al. 2017). For example, the recent establishment of the invasive spotted lanternfly (Lycorma delicatula White) in both North America and other locations around the globe has motivated several analyses involving SDMs (Jung et al. 2017; Wakie et al. 2020) and evaluations of potential control measures (Leach et al. 2019). In these and other examples, climatic niche information has been used to project the potential future distribution of a species (Wakie et al. 2020).

For a more thorough assessment of the risk of dispersal to PAs, natural areas, and agricultural lands, potential conservation corridor managers can also consider (1) the distribution and connectivity of primary host species; (2) the spatial overlap between the predicted climatic niche and the distribution of potential host species (e.g., there are thought to be more than 60 for spotted lantern fly; Lee et al. 2019); (3) the life-history traits of the focal species (dispersal ability, reproductive cycle, and phenology; Muthukrishnan et al. 2018); (4) the potential for human transport (direct or indirect); and (5) the risk to threatened or commercially important species (Andersen et al. 2004). Spatially explicit predictive risk assessments grounded in an understanding of a species biology are often necessary to improve predictions and the effectiveness of management (e.g., hemlock wooly adelgid (*Adelges tsugae* Annand) Liang et al. 2014, and Asian long-horned beetle (*Anoplophora glabripennis* Motschulsky; Favaro et al. 2015; Fig. 2, #2–4).

Scheduling and budgeting considerations

Protected area managers contend with an enormous array of management objectives (e.g., planning ecological restoration work, working with research scientists, planning an annual budget, or determining staffing requirements for a busy field season). In Canada, government funding for the management of PAs has historically been inadequate (Canadian Parks and Wilderness Society 2021) and has been documented at federal and provincial levels (Office of the Auditor General of Canada 2008; Office of the Auditor General of Ontario 2020). More recently, there has been a surge in government funding for conservation science and management to support Canada's adoption of the 2030 Global Biodiversity Framework from the United Nations Convention on Biological Diversity, which includes targets for PA connectivity (Government of Canada 2021) and support for invasive species management (Invasive Species Centre 2023). This recent focus on the need to make progress in PA connectivity and invasive species management arrives at a critical time where many PA managers are contending with increasing anthropogenic pressures and climate change impacts.

Many decision-support tools have been developed for invasive species management that integrate biophysical parameters and management scenarios. Typically, these are speciesspecific endeavours that integrate rates of spread and phenology of a given invasive species and also often include budgeting and scheduling scenarios to determine optimal longterm management strategies (Adams and Setterfield 2015). However, such approaches rarely consider the "connectivity conundrum" (Ashton et al. 2020; Saffariha et al. 2023, but see Minor and Gardner 2011). These modelling exercises focus on determining optimal timing and extent of ecological monitoring (Bonneau et al. 2018), expenditure on efforts focused on eradication (Adams and Setterfield 2015), and generally assessing the cost of labour and other resources required to carry out the work within a management cycle (Baker et al. 2017). In many cases, these integrative models are complex and may be difficult to implement, so a "science-practice gap" remains (Thompson et al. 2021). Developing more useful decision support tools may require a more collaborative approach that directly involves managers at various stages of the development process (Bodner et al. 2021). Collaborative development can take time, but it can also yield significant returns on investment for evidence-based management strategies (Hanley and Roberts 2019).

Partnerships and outreach

Recent analysis has revealed that the vast majority of PAs lack effective approaches for prioritizing the management of invasive species (Forner et al. 2022). Further, it is rarely the case that invasive species management plans are assembled using input from multiple stakeholders and rights holders to determine conservation priorities (Shackleton et al. 2019). Given the complex, multi-scale nature of biological invasions, effective prioritization of monitoring efforts and/or the application of control measures should benefit from extensive public engagement and collaboration (Crowley et al. 2017). There are examples of management frameworks that include stakeholder input, public perception, and expert opinion, where these considerations are integrated to form consensus opinion on the potential impact of an invader, specifically their potential impacts on biodiversity, ecosystem services, and public safety (Gaertner et al. 2017; Van Poorten and Beck 2021). Potgieter et al. (2018) developed a decision support model for the management of invasive species in Cape Town, South Africa, where these impact factors were assigned weighted values following consultation with stakeholder input and expert opinion. Outputs from this model included a spatially explicit characterization of invasion risk, site sensitivity, and optimal management action. Depending on the scale of PA connectivity planning, a decision support model may need to be structured in a way that integrates multiple actors to appropriately gauge both risk and effective management actions (Fig. 2, #5). In this regard, a generalized decision support framework that synthesizes considerations for both connectivity for conservation and invasive species management would be a welcome addition to the next national-scale biodiversity framework.

It is often the case that human movement into and within PAs acts as a primary driver of biological invasions (Guimarães Silva et al. 2020). As such, many ecosystem managers around the world have sought to improve their communications and public engagement methods in hopes of minimizing this invasion pathway (Lukács and Valkó 2021). These types of initiatives often involve a combination of zoning and PA signage, where human movement is regulated within a PA (Vardarman et al. 2018). Other strategies include engaging the broader public in the region of the PA to incentivize invasive species management (Drescher et al. 2019), accessing community science for early detection (Bonnet et al. 2020), and educational programming (Bravo-Vargas et al. 2019). Public engagement and inclusive approaches to PA management are now recognized as vital components of effective invasive species management practices in and around PAs (Shackleton et al. 2019). This is also true for PA connectivity planning, where community science, private land restoration, and adherence to planning policies all require buy-in from the public to improve the probability of success (Ban et al. 2013). Community science can also provide effective low-cost ecological monitoring in and around PAs (Binley et al. 2021) and contribute to regional PA connectivity planning to mitigate the potential for invasive species to disperse between PAs.

Concluding remarks

In addition to growing national commitments for protected area expansion, effective planning for the connectivity of PA networks and subsequent implementation will be critical in addressing dynamic ecological change in the coming decades (Hilty et al. 2020; UN Environment Program 2022). Central to these PA connectivity considerations (e.g., Targets 2 and 3) are the current and future spread and impacts of invasive species on native ecosystems and PAs (Shulze et al. 2018). The need to facilitate the movement of some species and impede others leads to connectivity conundrums. In Canada, this is of particular concern in the south of the country, where landscapes are highly fragmented and there is a disproportionate concentration of species of conservation concern (Kraus and Hebb 2020). We propose an integrative adaptive management framework and highlight actions that can help address these conundrums in short- and longterm PA planning. Hence, to improve their resistance to invasion, the management and design of PA networks need to focus not only on corridors but also on the PAs themselves as well.

Central to our perspective is the need for collaborative adaptive management for landscape connectivity considerations in PA network planning. However, we also acknowledge that PA managers are often resource-limited and, in many cases, are operating in a state of conservation triage (Coad et al. 2019; Dietz et al. 2021). Robust ecological monitoring and proactive risk assessments can inform both prospects for improved connectivity as well as the threat of species invasions. Further, the complexity of spatial planning for PA networks calls for effective engagement with regional partners (e.g., managers, land owners, Indigenous communities, scientists, and government agencies) and PA patrons. Anthropogenic disturbances, including human movement, act as both the drivers of biological invasions and impediments to conservation connectivity. Protected area planners can address these dynamic aspects of the connectivity conundrum through collaborative and integrative adaptive management planning.

Acknowledgements

This project was undertaken with the financial support of the Government of Canada through the federal Department of Environment and Climate Change. M-JF is thankful for an NSERC of Canada Discovery Grant (#5134) and the NSERC Canada Research Chair (CRC). We would also like to acknowledge the helpful feedback we received for this manuscript from one anonymous reviewer and our colleagues at the University of Toronto and Environment and Climate Change Canada.

Article information

Editor John P. Smol

History dates

Received: 26 June 2023 Accepted: 17 October 2023 Version of record online: 14 December 2023

Copyright

© 2023 The Author(s). This work is licensed under a Creative Commons Attribution 4.0 International License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Data availability

All relevant data are within the paper.

Author information

Author ORCIDs

Stuart W. Livingstone https://orcid.org/0000-0003-1031-8904 Josie Hughes https://orcid.org/0000-0001-7875-9015 Richard Pither https://orcid.org/0000-0002-1138-9408 Marie-Josée Fortin https://orcid.org/0000-0002-9935-1366

Author contributions

Conceptualization: SWL, JH, RP, MF Funding acquisition: JH, RP, MF Project administration: SWL, MF Supervision: JH, RP Writing – original draft: SWL, MF Writing – review & editing: SWL, JH, RP, MF

Competing interests

The authors declare there are no competing interests.

References

- Adams, V.M., and Setterfield, S.A. 2015. Optimal dynamic control of invasions: applying a systematic conservation approach. Ecological Applications, 25(4): 1131–1141. doi:10.1890/14-1062.1. PMID: 26465047.
- Andersen, M.C., Adams, H., Hope, B., and Powell, M. 2004. Risk analysis for invasive species: general framework and research needs. Risk Analysis: An Official Publication of the Society for Risk Analysis, 24(4): 893–900. doi:10.1111/j.0272-4332.2004.00487.x. PMID: 15357808.
- Ashton, I., Symstad, A., Baldwin, H., Post van der Burg, M., Bekedam, S., Borgman, E., et al. 2020. A new decision support tool for collaborative adaptive vegetation management in northern Great Plains national parks. Parks Stewardship Forum, 36(3): 510–518. doi:10.5070/ P536349865.
- Austin, M. 2007. Species distribution models and ecological theory: a critical assessment and some possible new approaches. Ecological Modelling, 200(1): 1–19. doi:10.1016/j.ecolmodel.2006.07.005.
- Baker, C.M., Armsworth, P.R., and Lenhart, S.M. 2017. Handling overheads: optimal multi-method invasive species control. Theoretical Ecology, 10(4): 493–501. doi:10.1007/s12080-017-0344-1.
- Ban, N.C., Mills, M., Tam, J., Hicks, C.C., Klain, S., Stoeckl, N., et al. 2013. A social–ecological approach to conservation planning: embedding social considerations. Frontiers in Ecology and the Environment, 11(4): 194–202. doi:10.1890/110205.
- Beaury, E.M., Finn, J.T., Corbin, J.D., Barr, V., and Bradley, B.A. 2019. Biotic resistance to invasion is ubiquitous across ecosystems of the United States. Ecology Letters, 23(3): 476–482. doi:10.1111/ele.13446. PMID: 31875651.
- Beazley, K.F., Hum, J.D., and Lemieux, C.J. 2023. Enabling a National Program for Ecological Corridors in Canada in support of biodiver-

sity conservation, climate change adaptation, and Indigenous leadership. Biological Conservation, **286**: 110286. doi:10.1016/j.biocon. 2023.110286.

- Beger, M., Metaxas, A., Balbar, A.C., McGowan, J.A., Daigle, R., Kuempel, C.D., et al. 2022. Demystifying ecological connectivity for actionable spatial conservation planning. Trends in Ecology & Evolution, 37(12): 1079–1091. doi:10.1016/j.tree.2022.09.002.
- Belote, R.T., Dietz, M.S., Jenkins, C.N., McKinley, P.S., Irwin, G.H., Fullman, T.J., et al. 2017. Wild, connected, and diverse: building a more resilient system of protected areas. Ecological Applications, 27(4): 1050–1056. doi:10.1002/eap.1527. PMID: 28263450.
- Beyer, H.L., Venter, O., Grantham, H.S., and Watson, J.E.M. 2019. Substantial losses in ecoregion intactness highlight urgency of globally coordinated action. Conservation Letters, 13(2): e12592. doi:10.1111/conl. 12692.
- Binley, A.D., Proctor, C.A., Pither, R., Davis, S.A., and Bennett, J.R. 2021. The unrealized potential of community science to support research on the resilience of protected areas. Conservation Science and Practice, 3(5): e376. doi:10.1111/csp2.376.
- Bixler, R.P., Johnson, S., Emerson, K., Nabatchi, T., Reuling, M., Curtin, C., et al. 2016. Networks and landscapes: a framework for setting goals and evaluating performance at the large landscape scale. Frontiers in Ecology and the Environment, 14(3): 145–153. doi:10.1002/fee.1250.
- Blossey, B., Endriss, S.B., Casagrande, R., Häfliger, P., Hinz, H., Dávalos, A., et al. 2020. When misconceptions impede best practices: evidence supports biological control of invasive Phragmites. Biological Invasions, 22(3): 873–883. doi:10.1007/s10530-019-02166-8.
- Bodner, K., Rauen Firkowski, C., Bennett, J.R., Brookson, C., Dietze, M., Green, S., et al. 2021. Bridging the divide between ecological forecasts and environmental decision making. Ecosphere, 12(12): e03869. doi:10.1002/ecs2.3869.
- Bonneau, M., Hauser, C.E., Williams, N.S.G., and Cousens, R.D. 2018. Optimal schedule for monitoring a plant incursion when detection and treatment success vary over time. Biological Invasions, 20(3): 741– 756. doi:10.1007/s10530-017-1572-4.
- Bonnet, P., Joly, A., Faton, J.-M., Brown, S., Kimiti, D., Deneu, B., et al. 2020. How citizen scientists contribute to monitor protected areas thanks to automatic plant identification tools. Ecological Solutions and Evidence, 1(2): e12023. doi:10.1002/2688-8319.12023.
- Booy, O., Mill, A.C., Roy, H.E., Hiley, A., Moore, N., Robertson, P., et al. 2017. Risk management to prioritise the eradication of new and emerging invasive non-native species. Biological Invasions, **19**(8): 2401–2417. doi:10.1007/s10530-017-1451-z.
- Bravo-Vargas, V., García, R.A., Pizarro, J.C., and Pauchard, A. 2019. Do people care about pine invasions? Visitor perceptions and willingness to pay for pine control in a protected area. Journal of Environmental Management, **229**: 57–66. doi:10.1016/j.jenvman.2018.07.018.
- Cadotte, M.W., Yasui, S.L.E., Livingstone, S., and MacIvor, J.S. 2017. Are urban systems beneficial, detrimental, or indifferent for biological invasion? Biological Invasions, 1–15. doi:10.1007/s10530-017-1586-y.
- Canadian Council on Invasive Species. 2023, February 3. *Home*. Canadian Council on Invasive Species. Available from https://canadainvasives. ca/
- Canadian Parks and Wilderness Society. 2021. The grades are in: a report on Canada's progress on protecting its land and oceans. Canadian Parks and Wilderness Society. Available from https://cpaws.org/wp-c ontent/uploads/2021/06/cpaws-reportcard2021-web.pdf.
- Chapman, D.S., Gunn, I.D.M., Pringle, H.E.K., Siriwardena, G.M., Taylor, P., Thackeray, S.J., et al. 2020. Invasion of freshwater ecosystems is promoted by network connectivity to hotspots of human activity. Global Ecology and Biogeography, 29(4): 645–655. doi:10.1111/geb. 13051.
- Clark, G.F., and Johnston, E.L. 2011. Temporal change in the diversity– invasibility relationship in the presence of a disturbance regime. Ecology Letters, 14(1): 52–57. doi:10.1111/j.1461-0248.2010.01550.x.
- Coad, L., Watson, J.E., Geldmann, J., Burgess, N.D., Leverington, F., Hockings, M., et al. 2019. Widespread shortfalls in protected area resourcing undermine efforts to conserve biodiversity. Frontiers in Ecology and the Environment, **17**(5): 259–264. doi:10.1002/fee.2042.
- Crowley, S.L., Hinchliffe, S., and McDonald, R.A. 2017. Invasive species management will benefit from social impact assessment. Journal of Applied Ecology, **54**(2): 351–357. doi:10.1111/1365-2664.12817.

- Cuddington, K., Sobek-Swant, S., Crosthwaite, J.C., Lyons, D.B., and Sinclair, B.J. 2018. Probability of emerald ash borer impact for Canadian cities and North America: A mechanistic model. Biological Invasions, 20(9): 2661–2677. doi:10.1007/s10530-018-1725-0.
- D'Aloia, C.C., Naujokaitis-Lewis, I., Blackford, C., Chu, C., Curtis, J.M.R., Darling, E., et al. 2019. Coupled networks of permanent protected areas and dynamic conservation areas for biodiversity conservation under climate change. Frontiers in Ecology and Evolution, **7**. Available from https://www.frontiersin.org/articles/10.3389/fevo.2019.00027.
- Dauer, J.T., and Jongejans, E. 2013. Elucidating the population dynamics of Japanese knotweed using integral projection models. PLoS One, 8(9). doi:10.1371/journal.pone.0075181.
- Decker, K.L., Pocewicz, A., Harju, S., Holloran, M., Fink, M.M., Toombs, T.P., and Johnston, D.B. 2017. Landscape disturbance models consistently explain variation in ecological integrity across large landscapes. Ecosphere, 8(4): e01775. doi:10.1002/ecs2.1775.
- Dickie, M., McNay, S.R., Sutherland, G.D., Cody, M., and Avgar, T. 2020. Corridors or risk? Movement along, and use of, linear features varies predictably among large mammal predator and prey species. Journal of Animal Ecology, **89**(2): 623–634. doi:10.1111/1365-2656.13130.
- Didham, R.K. 2010. Ecological consequences of habitat fragmentation. In Encyclopedia of life sciences. John Wiley & Sons. doi:10.1002/ 9780470015902.a0021904.
- Dietz, S., Beazley, K.F., Lemieux, C.J., St. Clair, C., Coristine, L., Higgs, E., et al. 2021. Emerging issues for protected and conserved areas in Canada. FACETS, **6**: 1892–1921. doi:10.1139/facets-2021-0072.
- Dillon, W.W., Lieurance, D., Hiatt, D.T., Clay, K., and Flory, S.L. 2018. Native and invasive woody species differentially respond to forest edges and forest successional age. Forests, **9**(7): 381. doi:10.3390/f9070381.
- Downey, P.O., Williams, M.C., Whiffen, L.K., Auld, B.A., Hamilton, M.A., Burley, A.L., and Turner, P.J. 2010. Managing alien plants for biodiversity outcomes—the need for triage. Invasive Plant Science and Management, 3(1): 1–11. doi:10.1614/JPSM-09-042.1.
- Drescher, M., Epstein, G.B., Warriner, G.K., and Rooney, R.C. 2019. An investigation of the effects of conservation incentive programs on management of invasive species by private landowners. Conservation Science and Practice, 1(7): e56. doi:10.1111/csp2.56.
- Drever, M.C., Smith, A.C., Venier, L.A., Sleep, D.J.H., and MacLean, D.A. 2018. Cross-scale effects of spruce budworm outbreaks on boreal warblers in eastern Canada. Ecology and Evolution, **8**(15): 7334–7345. doi:10.1002/ece3.4244.
- Emilson, C.E., and Stastny, M. 2019. A decision framework for Hemlock Woolly Adelgid management: review of the most suitable strategies and tactics for eastern Canada. Forest Ecology and Management, **444**: 327–343. doi:**10.1016/j.foreco.2019.04.056**.
- Environment and Climate Change Canada. 2010. Overabundant species: special conservation measures [Datasets; research]. Available from https://www.canada.ca/en/environment-climate-change/services/mig ratory-game-bird-hunting/overabundant-species-special-conservati on-measures.html.
- Environment and Climate Change Canada. 2016. 2020 Biodiversity goals & targets for Canada. Minister of Environment and Climate Change Canada. Available from https://publications.gc.ca/collections/collect ion_2016/eccc/CW66-524-2016-eng.pdf.
- Environment and Climate Change Canada. 2023. Toward a 2030 Biodiversity Strategy for Canada: halting and reversing nature loss. Environment and Climate Change Canada = Environnement et changement climatique Canada. Cat. No.: En4-539/2023E-PDF.
- Epanchin-Niell, R.S., Hufford, M.B., Aslan, C.E., Sexton, J.P., Port, J.D., and Waring, T.M. 2010. Controlling invasive species in complex social landscapes. Frontiers in Ecology and the Environment, **8**(4): 210–216. doi:10.1890/090029.
- Estévez, R.A., Anderson, C.B., Pizarro, J.C., and Burgman, M.A. 2015. Clarifying values, risk perceptions, and attitudes to resolve or avoid social conflicts in invasive species management. Conservation Biology, 29(1): 19–30. doi:10.1111/cobi.12359.
- Fantle-Lepczyk, J.E., Haubrock, P.J., Kramer, A.M., Cuthbert, R.N., Turbelin, A.J., Crystal-Ornelas, R., et al. 2022. Economic costs of biological invasions in the United States. Science of The Total Environment, 806: 151318. doi:10.1016/j.scitotenv.2021.151318.
- Favaro, R., Wichmann, L., Ravn, H.P., and Faccoli, M. 2015. Spatial spread and infestation risk assessment in the Asian longhorned beetle,

Anoplophora glabripennis. Entomologia Experimentalis et Applicata, **155**(2): 95–101. doi:**10.1111/eea.12292**.

- Forner, W.G., Zalba, S.M., and Guadagnin, D.L. 2022. Methods for prioritizing invasive plants in protected areas: a systematic review. Natural Areas Journal, 42(1): 69–78. doi:10.3375/20-47.
- Fournier, R.E., and Turgeon, J.J. 2017. Surveillance during monitoring phase of an eradication programme against *Anoplophora glabripennis* (Motschulsky) guided by a spatial decision support system. Biological Invasions, **19**(10): 3013–3035. doi:10.1007/ s10530-017-1505-2.
- Foxcroft, L.C., JaroŠÍK, V., Pyšek, P., Richardson, D.M., and Rouget, M. 2011. Protected-area boundaries as filters of plant invasions. Conservation Biology, 25(2): 400–405. doi:10.1111/j.1523-1739.2010. 01617.x.
- Foxcroft, L.C., Pyšek, P., Richardson, D.M., Genovesi, P., and Mac-Fadyen, S. 2017. Plant invasion science in protected areas: progress and priorities. Biological Invasions, 19: 1353–1378. doi:10.1007/ s10530-016-1367-z.
- Funk, J.L., Cleland, E.E., Suding, K.N., and Zavaleta, E.S. 2008. Restoration through reassembly: plant traits and invasion resistance. Trends in Ecology & Evolution, 23(12): 695–703. doi:10.1016/j.tree. 2008.07.013.
- Gaertner, M., Novoa, A., Fried, J., and Richardson, D.M. 2017. Managing invasive species in cities: a decision support framework applied to Cape Town. Biological Invasions, 19(12): 3707–3723. doi:10.1007/ s10530-017-1587-x.
- Giljohann, K.M., Hauser, C.E., Williams, N.S.G., and Moore, J.L. 2011. Optimizing invasive species control across space: willow invasion management in the Australian Alps. Journal of Applied Ecology, 48(5): 1286–1294. doi:10.1111/j.1365-2664.2011.02016.x.
- Government of Canada, D. of F. 2021, April 19. Budget 2021—Chapter 5: a healthy environment for a healthy economy. Available from https: //www.budget.canada.ca/2021/report-rapport/p2-en.html#chap5.
- Great Lakes Phragmites Collaborative. n.d. PAMF Technical Working Group | Great Lakes Phragmites Collaborative. Great Lakes Phragmites Collaborative. Available from https://www.greatlakesphragmit es.net/pamf/twg/ [accessed 26 February 2020].
- Guimarães Silva, R., Zenni, R.D., Rosse, V.P., Bastos, L.S., and van den Berg, E. 2020. Landscape-level determinants of the spread and impact of invasive grasses in protected areas. Biological Invasions, 22(10): 3083–3099. doi:10.1007/s10530-020-02307-4.
- Häkkilä, M., Abrego, N., Ovaskainen, O., and Mönkkönen, M. 2018. Habitat quality is more important than matrix quality for bird communities in protected areas. Ecology and Evolution, **8**(8): 4019–4030. doi:10.1002/ece3.3923.
- Hanley, N., and Roberts, M. 2019. The economic benefits of invasive species management. People and Nature, 1(2): 124–137. doi:10.1002/ pan3.31.
- Harper, K.A., and Macdonald, S.E. 2002. Structure and composition of edges next to regenerating clear-cuts in mixed-wood boreal forest. Journal of Vegetation Science, 13(4): 535–546. doi:10.1111/j. 1654-1103.2002.tb02080.x.
- Hierro, J.L., Villarreal, D., Eren, Ö., Graham, J.M., and Callaway, R.M. 2006. Disturbance facilitates invasion: the effects are stronger abroad than at home. The American Naturalist, 168(2): 144–156. doi:10.1086/ 505767.
- Hilty, J., Worboys, G.L., Keeley, A., Woodley, S., Lausche, B., Locke, H., et al. 2020. Guidelines for conserving connectivity through ecological networks and corridors. Best Practice Protected Area Guidelines Series No. 30. IUCN, Gland, Switzerland.
- Hirsh-Pearson, K., Johnson, C.J., Schuster, R., Wheate, R.D., and Venter, O. 2022. Canada's human footprint reveals large intact areas juxtaposed against areas under immense anthropogenic pressure. FACETS, 7: 398–419. doi:10.1139/facets-2021-0063.
- Hughes, J., Lucet, V., Barrett, G., Moran, S., Manseau, M., Martin, A.E., et al. 2023. Comparison and parallel implementation of alternative moving-window metrics of the connectivity of protected areas across large landscapes. Landscape Ecology, 38(6): 1411–1430. doi:10.1007/ s10980-023-01619-9.
- Invasive Species Centre. 2023, May 5. Ontario investing \$1 million to fight invasive species. Invasive Species Centre. Available from https://www.invasivespeciescentre.ca/ontario-investing-1-milli on-to-fight-invasive-species/.

- Jones, K.R., Venter, O., Fuller, R.A., Allan, J.R., Maxwell, S.L., Negret, P.J., and Watson, J.E.M. 2018. One-third of global protected land is under intense human pressure. Science, 360(6390): 788–791. doi:10.1126/ science.aap9565.
- Jung, J.-M., Jung, S., Byeon, D., and Lee, W.-H. 2017. Model-based prediction of potential distribution of the invasive insect pest, spotted lanternfly *Lycorma delicatula* (Hemiptera: Fulgoridae), by using CLIMEX. Journal of Asia-Pacific Biodiversity, **10**(4): 532–538. doi:10. **1016**/j.japb.2017.07.001.
- Kraus, D., and Hebb, A. 2020. Southern Canada's crisis ecoregions: identifying the most significant and threatened places for biodiversity conservation. Biodiversity and Conservation, 29(13): 3573–3590. doi:10. 1007/s10531-020-02038-x.
- Kullberg, P., Di Minin, E., and Moilanen, A. 2019. Using key biodiversity areas to guide effective expansion of the global protected area network. Global Ecology and Conservation, 20, e00768. doi:10.1016/ j.gecco.2019.e00768.
- Lamarre, J.-F., Legagneux, P., Gauthier, G., Reed, E.T., and Bêty, J. 2017. Predator-mediated negative effects of overabundant snow geese on arctic-nesting shorebirds. Ecosphere, 8(5): e01788. doi:10.1002/ecs2. 1788.
- Lawler, J.J., Wiersma, Y.F., and Huettmann, F. 2011. Using species distribution models for conservation planning and ecological forecasting. In Predictive species and habitat modeling in landscape ecology: concepts and applications. *Edited by* C.A. Drew, Y.F. Wiersma and F. Huettmann. Springer. pp. 271–290. doi:10.1007/978-1-4419-7390-0_ 14.
- Leach, H., Biddinger, D.J., Krawczyk, G., Smyers, E., and Urban, J.M. 2019. Evaluation of insecticides for control of the spotted lanternfly, *Ly-corma delicatula* (Hemiptera: Fulgoridae), a new pest of fruit in the Northeastern U.S. Crop Protection, **124**: 104833. doi:10.1016/j.cropro. 2019.05.027.
- Leberger, R., Rosa, I.M.D., Guerra, C.A., Wolf, F., and Pereira, H.M. 2020. Global patterns of forest loss across IUCN categories of protected areas. Biological Conservation, 241: 108299. doi:10.1016/j.biocon.2019. 108299.
- Lee, D.-H., Park, Y.-L., and Leskey, T.C. 2019. A review of biology and management of *Lycorma delicatula* (Hemiptera: Fulgoridae), an emerging global invasive species. Journal of Asia-Pacific Entomology, 22(2): 589–596. doi:10.1016/j.aspen.2019.03.004.
- Lemieux, C.J., Jacob, A., and Gray, P.A. 2021. Implementing connectivity conservation in Canada. Canadian Council on Ecological Areas. Available from https://ccea-ccae.org/wp-content/uploads/CCEA-Occasional Paper22-Connectivity-Low.pdf.
- Liang, J., He, X., Zeng, G., Zhong, M., Gao, X., Li, X., et al. 2018. Integrating priority areas and ecological corridors into national network for conservation planning in China. Science of the Total Environment, 626: 22–29. doi:10.1016/j.scitotenv.2018.01.086.
- Liang, L., Clark, J.T., Kong, N., Rieske, L.K., and Fei, S. 2014. Spatial analysis facilitates invasive species risk assessment. Forest Ecology and Management, 315: 22–29. doi:10.1016/j.foreco.2013.12.019.
- Lukács, K., and Valkó, O. 2021. Human-vectored seed dispersal as a threat to protected areas: prevention, mitigation and policy. Global Ecology and Conservation, 31: e01851. doi:10.1016/j.gecco.2021.e01851.
- MacDougall, A.S., Gilbert, B., and Levine, J.M. 2009. Plant invasions and the niche. Journal of Ecology, **97**(4): 609–615. doi:10.1111/j.1365-2745. 2009.01514.x.
- Maguire, D.Y., James, P.M.A., Buddle, C.M., and Bennett, E.M. 2015. Landscape connectivity and insect herbivory: a framework for understanding tradeoffs among ecosystem services. Global Ecology and Conservation, 4: 73–84. doi:10.1016/j.gecco.2015.05.006.
- Mansourian, S., Walters, G., and Gonzales, E. 2019. Identifying governance problems and solutions for forest landscape restoration in protected area landscapes. Parks, 25(1): 83–96. doi:10.2305/IUCN.CH. 2019.PARKS-25-1SM.en.
- Markle, C.E., Chow-Fraser, G., and Chow-Fraser, P. 2018. Long-term habitat changes in a protected area: implications for herpetofauna habitat management and restoration. PLoS One, **13**(2). doi:10.1371/journal. pone.0192134.
- Martin, P.H., and Marks, P.L. 2006. Intact forests provide only weak resistance to a shade-tolerant invasive Norway maple (*Acer platanoides* L.). Journal of Ecology, **94**(6): 1070–1079. doi:10.1111/j.1365-2745.2006. 01159.x.

- Matlack, G.R. 1993. Microenvironment variation within and among forest edge sites in the eastern United States. Biological Conservation, **66**(3): 185–194. doi:10.1016/0006-3207(93)90004-K.
- Meloche, C., and Murphy, S.D. 2006. Managing tree-of-heaven (Ailanthus altissima) in parks and protected areas: a case study of Rondeau Provincial Park (Ontario, Canada). Environmental Management, 37(6): 764–772. doi:10.1007/s00267-003-0151-x.
- Minor, E.S., and Gardner, R.H. 2011. Landscape connectivity and seed dispersal characteristics inform the best management strategy for exotic plants. Ecological Applications, **21**(3): 739–749. doi:10.1890/10-0321.1.
- Moodley, D., Angulo, E., Cuthbert, R.N., Leung, B., Turbelin, A., Novoa, A., et al. 2022. Surprisingly high economic costs of biological invasions in protected areas. Biological Invasions, **24**: 1995–2016. doi:10.1007/s10530-022-02732-7.
- Mortensen, D.A., Rauschert, E.S.J., Nord, A.N., and Jones, B.P. 2009. Forest roads facilitate the spread of invasive plants. Invasive Plant Science and Management, **2**(3): 191–199. doi:10.1614/IPSM-08-125.1.
- Mumma, M.A., Gillingham, M.P., Parker, K.L., Johnson, C.J., and Watters, M. 2018. Predation risk for boreal woodland caribou in humanmodified landscapes: evidence of wolf spatial responses independent of apparent competition. Biological Conservation, 228: 215–223. doi:10.1016/j.biocon.2018.09.015.
- Muthukrishnan, R., Davis, A.S., Jordan, N.R., and Forester, J.D. 2018. Invasion complexity at large spatial scales is an emergent property of interactions among landscape characteristics and invader traits. PLoS One, 13(5): e0195892. doi:10.1371/journal.pone. 0195892.
- Nackley, L.L., West, A.G., Skowno, A.L., and Bond, W.J. 2017. The nebulous ecology of native invasions. Trends in Ecology & Evolution, **32**(11): 814–824. doi:10.1016/j.tree.2017.08.003.
- Newbold, T., Hudson, L.N., Hill, S.L.L., Contu, S., Lysenko, I., Senior, R.A., et al. 2015. Global effects of land use on local terrestrial biodiversity. Nature, **520**(7545): 45–50. doi:10.1038/nature14324.
- Newmark, W.D., Halley, J.M., Beier, P., Cushman, S.A., McNeally, P.B., and Soulé, M.E. 2023. Enhanced regional connectivity between western North American national parks will increase persistence of mammal species diversity. Scientific Reports, 13(1): 474. doi:10.1038/ s41598-022-26428-z.
- Nobert, B.R., Merrill, E.H., Pybus, M.J., Bollinger, T.K., and Hwang, Y.T. 2016. Landscape connectivity predicts chronic wasting disease risk in Canada. Journal of Applied Ecology, **53**(5): 1450–1459. doi:10.1111/1365-2664.12677.
- Noss, R.F., and Harris, L.D. 1986. Nodes, networks, and MUMs: preserving diversity at all scales. Environmental Management, **10**(3): 299–309. doi:10.1007/BF01867252.
- Office of the Auditor General of Canada. 2008. 2008 Status Report of the Commissioner of the Environment and Sustainable Development to the House of Commons. Office of teh Auditor General of Canada. p. 28. Available from https://www.oag-bvg.gc.ca/internet/docs/aud_ch_c esd_200803_04_e.pdf.
- Office of the Auditor General of Ontario. 2020. Value-for-money audit: conserving the natural environment with protected areas, 2020 (p. 64). Office of the Auditor General of Ontario. Available from https://www.auditor.on.ca/en/content/annualreports/arreports /en20/ENV_conservingthenaturalenvironment_en20.pdf.
- Padmanaba, M., Tomlinson, K.W., Hughes, A.C., and Corlett, R.T. 2017. Alien plant invasions of protected areas in Java, Indonesia. Scientific Reports, 7(1): 1–11. doi:10.1038/s41598-017-09768-z.
- Parks Canada Agency. 2022. National program for ecological corridors nature and science. Available from https://parks.canada.ca/nature/sci ence/conservation/corridors-ecologiques-ecological-corridors.
- Perry, G.L.W., Moloney, K.A., and Etherington, T.R. 2017. Using network connectivity to prioritise sites for the control of invasive species. Journal of Applied Ecology, 54(4): 1238–1250. doi:10.1111/1365-2664. 12827.
- Pierre, S.M., Quintana-Ascencio, P.F., Boughton, E.H., and Jenkins, D.G. 2017. Dispersal and local environment affect the spread of an invasive apple snail (*Pomacea maculata*) in Florida, USA. Biological Invasions, 19(9): 2647–2661. doi:10.1007/s10530-017-1474-5.
- Pither, R., O'Brien, P., Brennan, A., Hirsh-Pearson, K., and Bowman, J. 2023. Predicting areas important for ecological connectivity through-

out Canada. PLoS One, **18**(2): e0281980. doi:10.1371/journal.pone. 0281980.

- Poley, L.G., Schuster, R., Smith, W., and Ray, J.C. 2022. Identifying differences in roadless areas in Canada based on global, national, and regional road datasets. Conservation Science and Practice, 4(4): e12656. doi:10.1111/csp2.12656.
- Potgieter, L.J., Gaertner, M., Irlich, U.M., O'Farrell, P.J., Stafford, L., Vogt, H., and Richardson, D.M. 2018. Managing urban plant invasions: a multi-criteria prioritization approach. Environmental Management, 62(6): 1168–1185. doi:10.1007/s00267-018-1088-4.
- Proctor, M.F., McLellan, B.N., Stenhouse, G.B., Mowat, G., Lamb, C.T., and Boyce, M.S. 2020. Effects of roads and motorized human access on grizzly bear populations in British Columbia and Alberta, Canada. Ursus, **2019**(30e2): 16. doi:10.2192/URSUS-D-18-00016.2.
- Pyšek, P., Jarošík, V., Hulme, P.E., Pergl, J., Hejda, M., Schaffner, U., and Vilà, M. 2012. A global assessment of invasive plant impacts on resident species, communities and ecosystems: the interaction of impact measures, invading species' traits and environment. Global Change Biology, 18(5): 1725–1737. doi:10.1111/j.1365-2486.2011. 02636.x.
- Quirion, B., Simek, Z., Dávalos, A., and Blossey, B. 2018. Management of invasive *Phragmites australis* in the Adirondacks: a cautionary tale about prospects of eradication. Biological Invasions, 20(1): 59–73. doi:10.1007/s10530-017-1513-2.
- Rejmánek, M., and Pitcairn, M.J. 2002. When is eradication of exotic pest plants a realistic goal. In Turning the tide: the eradication of invasive species. *Edited by* C. Veitch and M. Clout. IUCN. pp. 249–253.
- Resasco, J., Haddad, N.M., Orrock, J.L., Shoemaker, D., Brudvig, L.A., Damschen, E.I., et al. 2014. Landscape corridors can increase invasion by an exotic species and reduce diversity of native species. Ecology, 95(8): 2033–2039. doi:10.1890/14-0169.1.
- Richardson, D.M., Pyšek, P., Rejmánek, M., Barbour, M.G., Panetta, F.D., and West, C.J. 2000. Naturalization and invasion of alien plants: concepts and definitions. Diversity and Distributions, 6(2): 93–107. doi:10.1046/j.1472-4642.2000.00083.x.
- Riitters, K., Potter, K., Iannone, B.V., Oswalt, C., Fei, S., and Guo, Q. 2018. Landscape correlates of forest plant invasions: a high-resolution analysis across the eastern United States. Diversity and Distributions, 24(3): 274–284. doi:10.1111/ddi.12680.
- Rout, T.M., Hauser, C.E., McCarthy, M.A., and Moore, J.L. 2017. Adaptive management improves decisions about where to search for invasive species. Biological Conservation, 212: 249–255. doi:10.1016/j.biocon. 2017.04.009.
- Saffariha, M., Jahani, A., Roche, L.M., and Hosseinnejad, Z. 2023. Environmental decision support system development for natural distribution prediction of *Festuca ovina* in restoration of degraded lands. Land Degradation & Development, 1–20. doi:10.1002/ldr.4872.
- Sanderson, L.A., Mclaughlin, J.A., and Antunes, P.M. 2012. The last great forest: a review of the status of invasive species in the North American boreal forest. Forestry: An International Journal of Forest Research, 85(3): 329–340. doi:10.1093/forestry/cps033.
- Saura, S., Bertzky, B., Bastin, L., Battistella, L., Mandrici, A., and Dubois, G. 2019. Global trends in protected area connectivity from 2010 to 2018. Biological Conservation, 238: 108183. doi:10.1016/j.biocon.2019.07. 028.
- Saura, S., Bodin, Ö., and Fortin, M.-J. 2014. EDITOR'S CHOICE: stepping stones are crucial for species' long-distance dispersal and range expansion through habitat networks. Journal of Applied Ecology, 51(1): 171–182. doi:10.1111/1365-2664.12179.
- Schulze, K., Knights, K., Coad, L., Geldmann, J., Leverington, F., Eassom, A., et al. 2018. An assessment of threats to terrestrial protected areas. Conservation Letters, **11**(3): e12435. doi:10.1111/conl.12435.
- Sciance, M.B., Patrick, C.J., Weller, D.E., Williams, M.N., McCormick, M.K., and Hazelton, E.L.G. 2016. Local and regional disturbances associated with the invasion of Chesapeake Bay marshes by the common reed *Phragmites australis*. Biological Invasions, **18**(9): 2661–2677. doi:10.1007/s10530-016-1136-z.
- Senf, C., Campbell, E.M., Pflugmacher, D., Wulder, M.A., and Hostert, P. 2017. A multi-scale analysis of western spruce budworm outbreak dynamics. Landscape Ecology, **32**(3): 501–514. doi:10.1007/ s10980-016-0460-0.



- Shackleton, R.T., Adriaens, T., Brundu, G., Dehnen-Schmutz, K., Estévez, R.A., Fried, J., et al. 2019. Stakeholder engagement in the study and management of invasive alien species. Journal of Environmental Management, 229: 88–101. doi:10.1016/j.jenvman.2018.04.044.
- Silva, M., Massi, K., Negri, R., and Pedrosa, F. 2023. Jaguars and wild pigs indicate protected area connectivity in the south-east Atlantic Forest (Brazil). Environmental Conservation, 50(1): 22–30. doi:10.1017/ S03768929222000479.
- Simberloff, D., and Cox, J. 1987. Consequences and costs of conservation corridors. Conservation Biology, 1(1): 63–71. doi:10.1111/j.1523-1739. 1987.tb00010.x.
- Smith, A.L., Bazely, D.R., and Yan, N. 2014. Are legislative frameworks in Canada and Ontario up to the task of addressing invasive alien species? Biological Invasions, 16(7): 1325–1344. doi:10.1007/ s10530-013-0585-x.
- Sodhi, D.S., Livingstone, S.W., Carboni, M., and Cadotte, M.W. 2019. Plant invasion alters trait composition and diversity across habitats. Ecology and Evolution, **9**: 6199–6210. doi:10.1002/ece3.5130.
- Srivastava, V. 2019. Species distribution models (SDM): applications, benefits and challenges in invasive species management. CAB Reviews: Perspectives in Agriculture, Veterinary Science, Nutrition and Natural Resources, 14(020). doi:10.1079/PAVSNNR201914020.
- Stewart-Koster, B., Olden, J.D., and Johnson, P.T.J. 2015. Integrating landscape connectivity and habitat suitability to guide offensive and defensive invasive species management. Journal of Applied Ecology, 52(2): 366–378. doi:10.1111/1365-2664.12395.
- Stewart, F.E.C., Darlington, S., Volpe, J.P., McAdie, M., and Fisher, J.T. 2019. Corridors best facilitate functional connectivity across a protected area network. Scientific Reports, 9(1): Article 1. doi:10.1038/ s41598-019-47067-x.
- Sy, M., Keenleyside, K., Adare, K., Reader, B., Plante, M., and Deering, P. 2009. Protecting native biodiversity from high-impact invasive species through the protected areas of Parks Canada. Biodiversity, 10(2–3): 51–55. doi:10.1080/14888386.2009.9712843.
- Thompson, B.K., Olden, J.D., and Converse, S.J. 2021. Mechanistic invasive species management models and their application in conservation. Conservation Science and Practice, 3(11): e533. doi:10.1111/csp2. 533.
- Travers, E., Härdtle, W., and Matthies, D. 2021. Corridors as a tool for linking habitats—shortcomings and perspectives for plant conservation. Journal for Nature Conservation, 60: 125974. doi:10.1016/j.jnc. 2021.125974.
- Trombulak, S.C., and Frissell, C.A. 2000. Review of ecological effects of roads on terrestrial and aquatic communities. Conservation Biology, **14**(1): 18–30. doi:10.1046/j.1523-1739.2000.99084.x.
- UN Environment Program. 2022. Kunming-Montreal Global Biodiversity Framework. Convention on Biological Diversity. p. 14. Available from https://www.cbd.int/doc/c/e6d3/cd1d/daf663719a03902a9b116c 34/cop-15-l-25-en.pdf.

- Urziceanu, M.M., Cîşlariu, A.G., Nagodă, E., Nicolin, A.L., Măntoiu, D.S., and Anastasiu, P. 2022. Assessing the invasion risk of humulus scandens using ensemble species distribution modeling and habitat connectivity analysis. Plants 11(7): 857. doi:10.3390/plants11070857.
- Václavík, T., Kanaskie, A., Hansen, E.M., Ohmann, J.L., and Meentemeyer, R.K. 2010. Predicting potential and actual distribution of sudden oak death in Oregon: prioritizing landscape contexts for early detection and eradication of disease outbreaks. Forest Ecology and Management, 260(6): 1026–1035. doi:10.1016/j.foreco.2010.06.026.
- Van Poorten, B., and Beck, M. 2021. Getting to a decision: using structured decision-making to gain consensus on approaches to invasive species control. Management of Biological Invasions, 12(1): 25–48. doi:10.3391/mbi.2021.12.1.03.
- Vardarman, J., Berchová-Bímová, K., and Pěknicová, J. 2018. The role of protected area zoning in invasive plant management. Biodiversity and Conservation, 27(8): 1811–1829. doi:10.1007/s10531-018-1508-z.
- Wakie, T.T., Neven, L.G., Yee, W.L., and Lu, Z. 2020. The establishment risk of *Lycorma delicatula* (Hemiptera: Fulgoridae) in the United States and globally. Journal of Economic Entomology, **113**(1): 306–314. doi:10. 1093/jee/toz259.
- Wallace, R.D., Bargeron, C.T., and Reaser, J.K. 2020. Enabling decisions that make a difference: guidance for improving access to and analysis of invasive species information. Biological Invasions, **22**(1): 37–45. doi:10.1007/s10530-019-02142-2.
- Whittington, J., Hebblewhite, M., Baron, R.W., Ford, A.T., and Paczkowski, J. 2022. Towns and trails drive carnivore movement behaviour, resource selection, and connectivity. Movement Ecology, 10(1): 17. doi:10.1186/s40462-022-00318-5.
- Wilkerson, M.L. 2013. Invasive plants in conservation linkages: a conceptual model that addresses an underappreciated conservation issue. Ecography, 36(12): 1319–1330. doi:10.1111/j.1600-0587.2013. 00182.x.
- Wilson, M.C., Chen, X.-Y., Corlett, R.T., Didham, R.K., Ding, P., Holt, R.D., et al. 2016. Habitat fragmentation and biodiversity conservation: key findings and future challenges. Landscape Ecology, 31(2): 219–227. doi:10.1007/s10980-015-0312-3.
- With, K.A. 2002. The landscape ecology of invasive spread. Conservation Biology, 16(5): 1192–1203. doi:10.1046/j.1523-1739.2002.01064.x.
- Wright, P., Lazaruk, H., Bjorgan, L., Gonzales, E., and Thurston, E. 2017. Discussion paper: evaluating management effectiveness (Pathway to Canada Target 1: Expert Task Team). p. 101.
- Yannelli, F.A., Hughes, P., and Kollmann, J. 2017. Preventing plant invasions at early stages of revegetation: the role of limiting similarity in seed size and seed density. Ecological Engineering, **100**: 286–290. doi:10.1016/j.ecoleng.2016.12.001.
- Yu, D., Xun, B., Shi, P., Shao, H., and Liu, Y. 2012. Ecological restoration planning based on connectivity in an urban area. Ecological Engineering, 46: 24–33. doi:10.1016/j.ecoleng.2012.04.033.